

ORIGINAL ARTICLE

Identification of Elite Cultivars in Maize (*Zea mays* L.) Germplasm Based on Agronomical CharactersF. J. B. Quenum¹, M. C. Djaboutou², S. S. Houedjissin^{3*}, M. G. Sinha², C. Houenou¹, G. H. Cacaï³, C. Ahanhanzo³¹Department of Plant Production, Faculty of Agronomic Sciences, University of Abomey-Calavi, Abomey Calavi, Benin,AQ1 ²Department of ???, National Institute of Agricultural Research of Benin, Center for Agricultural Research Cotton and Fibers, Benin, ³Department of Genetic and Biotechnologies, Faculty of Science and Technology, Abomey-Calavi University, Benin

ABSTRACT

AQ2 Morphological characterization is a first approach to diversity study genetic. 10 corn accessions collected in Katagon regions Zoungbomin and Ifangni of South Benin were evaluated morphologically in a device in randomized blocks. 24 morphological characters selected from maize descriptors were used in this study. The descriptive analysis revealed a significant diversity among accessions, marked by a significant gap between the minimum and maximum values. Thus, eight variables (male flowering to 50%; 50% female flowering, plant height, height of ear insertion, number of primary branch of the tassel, average number of kernels per row, weight of 100 grains, maturity), contributing most discriminating varieties Lagos Carder despite their performance. These characters can be basic criteria for identifying maize varieties in Benin.

Key words: Benin, genetic diversity, maize, morphological, variety

INTRODUCTION

Maize (*Zea mays* L.) is the largest crop in the world and the first cereal produced front of wheat, with a production of 843 million tons in 2014, against 653 million tons for wheat.^[1] In Benin corn, is also the most cultivated cereal, and holds first place with nearly 70% of cereal area planted. In southern Benin, it is the staple food with a consumption of forms of diversity (fresh corn, roasted corn, corn dough patty). Its production is estimated at over 900,000 tons, accounting for about four-fifths of the national cereal production. Despite the favorable conditions enjoyed by the culture, he knows an evolution saw tooth;^[2] although the research conducted in recent years have led to the development of improved varieties, some of which are already popular;^[3] it must be noted that these improved varieties are very few adopted by the producers for several reasons including their requirements in specific input and technological and organoleptic qualities that do not always meet the needs of users.^[4-6] Thus, there arises the need to highlight some local maize varieties that stand out from other varieties and meet the requirements of producers and consumers. Moreover, in the current highly unstable agroclimatic conditions, it is

urgent to screen varieties/ecotypes to highlight the best performers. This study aims to evaluate the performance of 10 varieties of maize in agroecological conditions of South Benin, to identify the best.

Study Site

The experiment was conducted from 29 May to 12 September 2014 at the experimental site of Faculty of Agricultural Sciences of the UAC to Sekou (town of Allada) (2°19'E, 6°12'N), located at 50 km north of Cotonou. The climate is subequatorial with two rainy seasons and two dry seasons. There is an average 1200 mm of water for about 85 days of rain per year. The average temperature varies between 27 and 29°C with relative humidity averaged 70%. The vegetation is of Guinean. The soil, commonly known as earth bar, is ferralitique kind desaturated to kaolinite.

MATERIAL AND METHODS

Plant Material

The plant material used as part of our work is exclusively maize. Table 1 present the 10 local varieties used and their characteristics.

Methods

Sampling design

The experimental design is a randomized block Fisher with four repetitions, of which 10 treatments

*Corresponding author:
Email: sergesth01@yahoo.frAQ3 Received: ???, Accepted: ???,
Published: ***http://dx.doi.org/***
ISSN *** © 2017

(10 varieties of corn) namely Massahouégotinwéwé, Massahouégotivé, Lagos, Carder, Massahouéwéwé, Massahoué, Massahouéwéwé, Massahouéwéwé, Massahouéwéwé, Massahouévèwè, used in this study. Land dimensions are 3-1.2 m. The spacing between planting holes is 0.4 m × 0.8 m and the spacing between lines is 0.8 m. **He alleys between elementary repetitions and pieces are, respectively, 1 and 0.6 m.**

AQ5

Cultural operations, measured parameters, and data collected

The test site was mowed and plowed flat, then fertilized by the intake of 78 g a mixture of triple super phosphate and 98 g K₂O per basic plot as basal dressing. The sowing took place after soil preparation and basal dressing at 2 grains by seed hole. The first weeding and hoeing the second held, respectively, 15 and 45 JAS JAS. The plants were thinned to 46 JAS stage five sheets; then a urea intake of 36 g by elemental plot was made. The harvest took place 107 days after sowing.

The following actions were taken: Plant height, length of plant, plant width, circumference collar, male flowering female flowering, branching away from the panicle, number of primary branch of the panicle, panicle length, number of ears by foot, ear insertion height, leaf number above the ear, spike length, diameter of the spike, plant height, circumference of the collar of the adventitious root, maturity, number of stored grain, average grain, weight of 100 grains, grain color, grain length, grain width, grain diameter, according to the descriptors of maize.

Statistical Analysis of Data

Statistics 8 software was used for analysis of variance (ANOVA) to compare means using the least square difference test at the 5% threshold. A descriptive statistical analysis was conducted to investigate the differences between the coefficients of variation for the characters semi-cycle of flowering and plant maturity, while the variance analysis focused on the characters number of primary branch of the panicle, number of ear per plant, height of insertion of ear and plant height. Minitab the software has achieved the matrix correlation.

RESULTS

Qualitative Character Analysis in Prospective Areas

The color of the grains presents a variation on all 10 varieties. White corn grains account for 50%. They are cultivated in the areas surveyed than yellow corn (40%) and red corn (10%) (Table 2).

The distribution of maize accessions according to color silk tassel and color shows through against 70% are green and 30% are purple (Table 3).

Descriptive Analysis of Quantitative Traits

Significant differences were observed between the minima and maxima for important agronomic traits such as male flowering (FLOM), silking (FLOF), the height of plant (HP),

AQ6

Table 1: Characteristics of varieties studied

Name of the cultivar	Cycle according to the producer	Ethnic group	Status of the sample	Sample origin	Frequency	Resistance to the dryness	Vermin resistance in stock	Color
Massahouégotinwéwé	2 months ½	Goun	Cultivar of the territory	Garret	Abundant	No	Average	Yellow
Massahouégotinvè	3 months ½	Goun	Cultivar of the territory	Garret	Abundant	Average	Average	Yellow
Lagos	3 months	Goun	Imported	Garret	Occasional	Average	Bad	Red
Cader	2 months ½	Goun	Improved	Garret	Occasional	Maid	Bad	White
Massahouéwéwé	2 months ½	Goun	Cultivar of the territory	Garret	Abundant	Maid	Maid	White
Massahoué	3 months ½	Goun	Cultivar of the territory	Garret	Abundant	Maid	Maid	Yellow
Massahouéwéwé	3 months	Goun	Cultivar of the territory	Garret	Abundant	Maid	Average	White
Massahouéwéwé	2 months ½	Nago	Cultivar of the territory	Garret	Frequent	Maid	Maid	White
Massahouéwéwé	2 months ½	Goun	Cultivar of the territory	Garret	Frequent	Maid	Maid	White
Massahouévèwè	3 months ½	Goun	Cultivar of the territory	Garret	Abundant	Maid	Average	White

Table 2: Distribution of grains color in the areas surveyed

Character analysis	Numbers of observation by area					
	Colors	Katagon	Zoungbomin	Ifangni centre	Total	Frequencies (%)
Grains color	Yellow	2	1	1	4	40
	Red	1	0	0	1	10
	White	2	1	2	5	50
	Total	5	2	3	10	100

number of grain rows (NRG) the average number of kernels per row (NMGR), weight of 100 grains (P100G) (Table 4). On average, accessions have a 50-day semi-flowering cycle. The earliest began his male flowering from 47 days after sowing and later to 52 days. Silking varies from 51 days to the earliest and 55 days later. The plant height varies from 131.65 cm for small varieties to 149.35 cm for tall varieties, with an average of 142.05 cm. The average number of kernels per row is 20 for accessions with small ears and 26 accessions large ears, with an average of 23.980. The weight of 100 grains varies from 21.075 g for accessions with small grains to 28,300 g for those coarse.

Table 5 obtained after the ANOVA, it appears that Massahouéwéwé over primary branch (20.57), Massahoué has the number of the lowest primary branch (13.5). Massahouégontinwéwé and Massahouégontivé (2.250) have more ears while Massahouévèwè (1.250) fewer ears. Regarding the insertion height of the ears

Massahouégontivé has the greatest height (87.50 cm) and the lowest Massahouéwéwé height of insertion of the ears (62 cm), with a significant difference at alpha 5% threshold.

Structuring of Morphological Diversity

A significant correlation (≥ 0.50) was observed between several pairs of variables (Table 6). Positive relationship between the circumference of the collar, branching away from the tassel, height insertion ears, the maturity and the number of row of grain with height plant which implies that an increase in plant height causes an increase of all these variables. Hence, the tall cultivars are late, have a good ear insertion height and a good circumference collar.

Analyze Differences between Flowerings

From descriptive analysis (Table 7), the coefficient of variation is low (<15%) for most of the measured variables

Table 3: Distribution of color tassel and silking

Character analysis	Color	Treatments										Total	Frequency
		T1	T2	T3	T4	T5	T6	T7	T8	T9	T10		
Panicle color	Green	4	42	2	2	2	2	3	3	3	3	28	70
	Purple	0	0	2	2	2	2	1	1	1	1	12	30
	Total	4	4	4	4	4	4	4	4	4	4	40	100
Silk color	Green	4	4	2	2	2	2	3	3	3	3	28	70
	Purple	0	0	2	2	2	2	1	1	1	1	12	30
	Total	4	4	4	4	4	4	4	4	4	4	40	100

AQ9

Table 4: Minimum, maximum values, mean, SD, CV of 19 quantitative traits studied

Variables	\bar{X}	Max	Min	SD	CV (%)
Circumference (cm)	5.6453	6.3000	5.0700	0.5876	20.82
L (cm)	59.342	69.410	46.155	8.9551	30.18
L (cm)	21.420	24.070	18.085	14.790	138.10
FLOM (jrs)	49.625	52.000	47.750	2.5228	10.17
FLOF (jrs)	52.800	55.000	51.250	2.42447	9.18
DRP (cm)	11.805	13.065	10.560	1.0272	17.40
LP (cm)	33.195	35.270	31.395	1.7938	10.81
HPL (cm)	142.05	149.95	131.61	8.4042	11.83
NFE	5.6700	6.0500	5.4000	0.2559	9.03
CAR (cm)	5.5235	5.8700	4.9850	0.2565	9.29
Mat (jrs)	115.83	99.00	91.00	63.903	110.34
LE (cm)	107.68	121.35	96.51	8.8042	16.35
DE (cm)	34.590	35.859	32.204	2.3085	13.35
NGR	11.790	12.700	11.150	0.7082	12,01
NMGR	23.980	26.650	20.600	2.4079	20.08
P100G (g)	24.815	28.300	21.075	2.4619	19.84
DG (cm)	4.5242	4.7100	4.1985	0.4263	18.84
LG (mm)	9.3027	9.5900	8.5845	0.5276	11.34
IG (mm)	8.6004	8.9910	7.8650	0.4368	10.16

\bar{X} : Average, Max: Maximum value, Min: Minimal value, SD: Standard of deviation, CV: Coefficient of variation, FLOM: Male flowering, FLOF: Silking, HP: Height of plant, NRG: Number of grain rows, NMGR: The average number of kernels per row, P100: Weight of 100 grains

which confirms the precision of the test. Furthermore, significant differences were recorded between the averages for some varieties. The differences in the days of the appearance of flowering male and female flowering is 2.75 days for varieties Lagos Carder with respective maturities of 91 and 90 days. The coefficients of variation are <10% for the three traits studied.

Analysis of Yield Components Varieties Lagos and Carder

Table 8 presented the coefficients of variation of some performance parameters for varieties Lagos Carder. It

appears from this table that the average values of all the parameters vary very little from one variety to the other with the exception of the length of the ear, where there is a significant difference of 10.13 mm. The coefficient of variation is low (<5%) for all variables.

Variations Height of the Various Varieties during the Test

The differences between the heights of the varieties Lagos and Carder vary from 1.77 cm to 9.36 cm, respectively, from 25 to 70 JAS (Figure 1).

Table 5: Change in the number of primary branches, number of cob by foot and height insertion ears

Treatment	Numbers of primary branch of the panicle	Number of ears per seedling	Height of insertion of ears (cm)
T1	18.650 ^{ab}	2.2500 ^a	81.000 ^{ab}
T2	16.050 ^{ab}	2.2500 ^a	87.500 ^a
T3	15.300 ^{ab}	1.7500 ^{abc}	78.050 ^{ab}
T4	15.450 ^{ab}	1.5000 ^{bc}	71.700 ^{ab}
T5	14.850 ^{ab}	2.0000 ^{ab}	75.850 ^{ab}
T6	13.500 ^b	1.7500 ^{abc}	70.380 ^{ab}
T7	20.575 ^a	2.0000 ^{ab}	62.000 ^b
T8	17.400 ^{ab}	2.0000 ^{ab}	72.600 ^{ab}
T9	15.200 ^{ab}	2.0000 ^{ab}	72.600 ^{ab}
T10	18.125 ^{ab}	1.2500 ^c	72.950 ^{ab}
P	0.03*	0.0265*	0.022*
LSD _{0.05}	6.9761	0.5746	24.656

AQ10 ^{a,b,c: ???}, LSD: Least significant difference

Table 6: Correlation matrix of the studied characters

	HP	CirC	LF	IF	FLOM	FLOF	DRP	LP	NBP	NEP	HIE	NFE	Mat	LE	DE	NGR	NMGR	P100	LG	IG	DG	
HP	1																					
CirC	0.68	1																				
LF	0.5	0.31	1																			
Lf	-0.2	-0.1	0.41	1																		
FLOM	0.13	-0.1	-0.5	-0.2	1																	
FLOF	0.05	-0.1	-0.6	-0.2	0.97	1																
DRP	0.83	0.4	0.51	-0.1	0.39	0.23	1															
LP	0.49	0.76	0.35	-0.3	-0.1	-0.3	0.47	1														
NBP	-0.1	-0.1	-0.3	0.12	0.36	0.34	0.2	-0.1	1													
NBP	0.23	0.02	0.01	-0.4	0.63	0.55	0.55	0.21	0.13	1												
HIE	0.89	0.5	0.34	-0.1	0.45	0.36	0.86	0.27	0.14	0.42	1											
NFE	0.12	0.21	0.72	0.19	-0.8	-0.9	0.05	0.45	-0.4	-0.4	-0.2	1										
Mat	0.51	0.49	0.14	0.13	0.59	0.56	0.5	0.13	-0.1	0.44	0.65	-0.4	1									
LE	0.26	0.22	0.47	-0.1	-0.4	-0.5	0.13	0.48	-0.7	-0.1	0.06	0.61	-0.2	1								
DE	0.16	0.34	0.57	0.31	-0.6	-0.6	-0.1	0.28	-0.7	-0.5	-0.1	0.67	0.06	0.63	1							
NGR	0.69	0.48	0.35	0.17	0.41	0.29	0.66	0.32	-0.2	0.29	0.83	-0.1	0.68	0.39	0.16	1						
NMGR	0.43	0.44	0.49	0.16	-0.0	-0.1	0.34	0.5	-0.6	0.06	0.34	0.37	0.42	0.78	0.68	0.73	1					
P100G	0.1	-0.2	0.25	-0.4	-0.5	-0.5	-0.0	-0.0	-0.2	-0.1	-0.2	0.46	-0.6	0.24	0.13	-0.48	-0.246	1				
LG	0.47	-0.1	0.27	-0.2	0.07	0.11	0.32	-0.3	-0.4	0	0.33	0.02	0.27	0.05	0.29	0.15	0.154	0.456	1			
IG	-0.5	-0.6	-0.2	-0.2	-0.2	-0.3	-0.4	-0.2	-0.1	-0.3	-0.6	0.31	-0.7	0.25	0.09	-0.54	-0.138	0.545	0.11	1		
DG	0.15	0.09	0.39	0.08	-0.4	-0.05	0.21	0.32	-0.1	-0.3	-0.1	0.73	-0.3	0.33	0.27	-0.01	0.213	0.226	0	0.41	1	

FLOM: Male flowering, FLOF: Silking, HP: Height of plant, NRG: Number of grain rows, NMGR: The average number of kernels per row, P100: Weight of 100 grains

Table 7: Change in flowering cycles and maturity

Treatment	FLOM		FLOF		Mat	
	$\bar{X} \pm SD$	CV	$\bar{X} \pm SD$	CV	$\bar{X} \pm SD$	CV
T1	52.00±6.48	12.46	55.00±6.480	11.78	99.00±8.08	8.16
T2	51.75±5.61	10.86	55.00±5.83	10.60	90.50±4.78	3.67
T3	47.75±4.50	9.42	50.50±4.50	8.92	91.00±2.70	2.97
T4	48.50±4.20	8.66	51.25±4.11	8.02	90.25±3.09	3.43
T5	49.25±4.27	8.67	52.25±3.50	6.69	99.00±8.36	8.45
T6	49.00±5.47	11.17	52.25±5.12	9.80	97.25±9.67	9.94
T7	49.25±4.57	9.28	52.25±4.50	8.61	93.25±8.53	9.15
T8	50.00±3.55	7.11	53.25±3.59	6.74	95.50±7.00	7.32
T9	49.25±4.57	9.28	53.00±4.54	8.57	95.00±7.78	8.19
T10	49.50±6.35	12.83	53.25±5.43	10.21	94.25±7.36	7.81

T1: Massahouégontinwéwé, T2: Massahouégontivé, T3: Lagos, T4: Carder, T5: Massahouéwéwé, T6: Massahoué, T7: Massahouéwéwé, T8: Massahouéwéwé, T9: Massahouéwéwé, T10: Massahouévèwè, SD: Standard of deviation, CV: Coefficient of variation, FLOM: Male flowering, FLOF: Silking

Table 8: Change in components of yield varieties Lagos and Carder

Treatment	P100G	LE	DE	NMGR	NRG
T3		121.35	35.587	26.40	12.20
T4		111.22	35.855	25.05	11.70
X	25	116.285	35.721	25.72	11.95
SD	0.575	5.065	0.134	0.675	0.25
CV	2.30	4.35	3.75	2.62	2.09

NRG: Number of grain rows, NMGR: The average number of kernels per row, P100: Weight of 100 grains, SD: Standard of deviation, CV: Coefficient of variation

Variations Height of the Varieties Lagos and Carder

Figure 2 presents the evolution height of the seedlings according to time. It arises that the difference between the Lagos varieties and to card is increasing 1.76, 4.08, 7.14, respectively, to 25, 35, 45 JAS. In addition, the difference in height observed with 70 JAS which is 2.40 cm show a reduction of the differences between height seedlings after flowering.

DISCUSSION

In the plan agromorphological, local accessions of corn grown in the Department of Ouémé in Benin revealed significant differences between the characters analyzed, indicating a strong phenotypic heterogeneity between 10 accessions. This morphological diversity was structured into six groups that differ in plant height, the semi-flowering cycle, the number of stored grain, the average number of kernels per row and weight of 100 grains. **These characters have helped distinguish the early and intermediate groups corn, midsize, late corn and groups large and imposing corn groups morphologically.** Similar results were obtained by Attiey^[7] in Ivory Coast and to suggest that morphological differentiation is often based on agronomic traits.^[8] The large differences between the minimum and maximum values for all analyzed morphological characters show significant inter-variability accessions. This strong result

morphological diversity of peasant seed management practices.^[9-12] The country phenotypical selection based on the characters perceptible (phenologic, vegetative, and ear) could explain the contribution of these variables to the structuring of variability.

A weak variation enters the averages of the varieties Carder and Lagos with a coefficient of variation lower than 10%. The transition from the male inflorescence to the female inflorescence occupies an interval of time from 2 to 4 days. These results are in conformity with those of Fleury,^[13] which supports that the necessary interval between male flowering and female flowering is lower than 5 days. The variations of days of appearance between male flowering and female flowering are 2.75 days for the varieties Lagos and Carder. More the shift between flowerings male and female is significant, less the material considered flowers in a synchronous way. This interval is influenced by the hydrous stress. The difference between flowerings male and female is very significant in improvement varietal because it directly influences the fecundation of the female flowers and thus the formation of grains. Moreover, the dryness is responsible for an increase in the difference between male and female flowering. Our results are similar to those of Sanou^[8] which undertook a study on the performance of the S2 lines of FBC6 compared to the hydrous stress. It observed a shift between flowerings male and female. On the other hand, our results are contrary with those of Zoma^[14] which undertook a study on the S1 lines of HOPE and showed that there is no significant difference between the lines compared to the shift between male and female flowering.

During this study, the varieties Lagos and Carder were identified by the variations reduced between the dates of male and female flowering, and their maturity rather short early (90 days). Indeed the test knew 32 days without rain, covering the significant stages of flowering.




Figure 1: Evolution the height seedling of the 10 varieties




Figure 2: Evolution height of the varieties Lagos and Carder

In spite of this condition of hydrous stress, the variations of days between the appearances of flowers male and female remained lower than 4.5 days for all the studied varieties, and 2.75 days for the varieties Lagos and Carder. This testifies to the aptitude of the varieties to tolerate the dryness in period of irregularity of the rains. Moreover, according to our investigations, the Carder variety would be initially variety enhanced vertical deepwater tree, introduced into the zone approximately 15 years ago.

CONCLUSION

The preliminary results on the diversity and the morphological characterization of corn cultivated in Ouémé clearly show that the different one are varieties tested present a variation for the whole of the characters used, in particular, those related to the cycle of semi-flowering, with the height of the plant, the number of lines of grain, the average number of grains per line and with the weight of 100 grains. This genetic variability observed between accessions constitutes an asset for work of selection. The differences observed between the varieties indicate that the varieties Lagos (P100G

24.42; NMGR 26.40; NRG 12.20; FLM 47.75; FLF 50.50) and Carder (P100G 25.57; NMGR 25.05; NRG 11.70; FLM 48.50; FLF 51.25), introduce no nearly but intermediate individuals. They are essential morphologiquement with ears inserted into low height (HIE Lagos 78.05 cm; to card 71.05 cm). The Lagos varieties and to Carder have the best characteristics of ear and grains and can be used as source of parents to improve the output. These two varieties are equipped with the best vegetative characteristics and could be proposed to the producers in these times of climatic changes.

REFERENCES

1. Louette, D. (1994). Traditional Maize Varieties in the Management of the Reserve Biosphere Sierra Manantan (RBSM, Jalisco and Colima States, Mexico) *In Situ* Conservation of Genetic Resources of Cultivated Plants. PhD Thesis. France: Agronomic School Montpellier. p245.
2. MAEP (Ministry of Agriculture, Livestock and Fisheries). (2013). Press Sheet management budget. Washington, DC: Ministry of Agriculture, Livestock and Fisheries. p1.
3. Yallou, C. G., Menkir, A., Adetimirin, V. O., & Kling, J. G. (2005). Combining Ability of diverse maize inbred lines under Striga infestation in Nigeria and Benin. In: Badu-Apraku, B.,

- Ouedraogo, M. A. B., Carsky, M., & Menkir, R. M. J. Fith Biennal West and Central Africa. Regional Maize Workshop. Benin: IITA. p12-27.
4. Kydd, J. (1989). Maize research in Malawi: Lessons from a failure. *Journal of International Development*, 1(1), 112-114. doi:10.1002/jid.3380010105.
 5. Koudokpon, V. (1991). Why the improved maize varieties are not adopted by farmers. *Agronomic Research Bulletin of Benin*, 2, 6-9.
 6. Tchamo, P. (1993). Maize improvement. Strategies intended for human consumption in eastern Cameroon. In: *Genetic Progress He Goes Through the Identification and Inventory of Genes? Paris-France: AUPELF-UREF, John Libbey Eurotext.* p347-357.
 7. Attiey, K. (1991). Contribution to the assessment of the variety of landraces of maize Ivory Coast Center. In: Menyonga, J. M., Bezuneh, T., Yayock, J. Y., & Sumana, I., editors. *Progress in the Search for Food Crop Production in Semi-Arid Africa. Scientific Communications Presented at the Inter-Network Conference of the OAU/STRC-SAFGRAD on Research and Production of Food Crops in Semi-Arid Regions of Sub-Saharan Africa, Niamey, Niger, 7-14 March; 1991.* p213-223.
 8. Sanou, A. (2011). Creation and evaluation of "hybrids and maize lines as part of" an intensification of maize farming in Burkina Faso. End of Cycle Memory. Bobo Dioulasso, Burkina Faso: IDR, UPB. p80.
 9. McKeye, D., Emperaie, L., Elias, M., Pinton, F., Robert, T., Desmoulière, S., & Rival, L. (2001). Regional and local dynamic management of varietal diversity of cassava in Amozonie. *Genetic Selection Evolution*, 33, 465-490.
 10. Barnaud, A., Joly, H. I., McKey, D., Deu, M., Khasah, C., Monné, S., & Garin, E. (2008). Management genetic resources of sorghum (*Sorghum bicolor*) in Duupa (North Cameroon). *Cahiers Agricultures*, 17(2), 178-182.
 11. Delaunay, S., Tesca, R. P., Oualbego, A., Vom Brocke, K., & Lançon, J. (2008). Cotton cultivation does not overturn the traditional trade of sorghum seed. *Books Agricultures*, 17, 189-194.
 12. Missihoun, A., Agbangla, C. A., Adoukonou-Sagbadja, H., Ahanhanzo, C., & Vodouhe, R. (2012). Traditional management and status of genetic resources of sorghum (*Sorghum bicolor* L. Moench) Northwest of Benin. *International Journal of Biological Chemical Sciences*, 6, 1003-1018. doi:10.4314/ijbcs.v6i3.8.
 13. Fleury. (2010). *Technical problems of sowing. French Association for the Study of the Sol. (2010). Available from: <http://www.afes.fr>. Food Biotechnology Council. (1990). Biotechnologies and food: Assuring the safety of foods produced by genetic modification. *Regulatory Toxicology and Pharmacology*, 12, S1-S196.*
 14. Zoma, W. O. (2010). Improved Maize Variety Hope for the Intensification of Its Culture, Dissertation Cycle. Bobo Dioulasso, Burkina Faso: IDR-UPB. p51.

AQ7,8

Author Queries???

AQ1: Kindly provide department

AQ2: Kindly provide structured abstract

AQ3: Kindly provide history details

AQ4: Kindly provide running title

AQ5: Kindly review the sentence.

AQ6: Kindly check the abbreviation and expansion

AQ7: Kindly provide complete reference details

AQ8: Please note three references seems to be combined as single reference. Kindly suggest whether to delete the latter or not.

AQ9: Kindly check the edit throughout the file.

AQ10: Kindly provide significant values